

Kochen-Specker theorem for 8-dimensional space

Michael Kernaghan

Department of Philosophy, University of Western Ontario, London N6A 3K7, Canada

and

Asher Peres

Department of Physics, Technion – Israel Institute of Technology, 32 000 Haifa, Israel

Abstract

A Kochen-Specker contradiction is produced with 36 vectors in a real 8-dimensional Hilbert space. These vectors can be combined into 30 distinct projection operators (14 of rank 2, and 16 of rank 1). A state-specific variant of this contradiction requires only 13 vectors, a remarkably low number for 8 dimensions.

The Kochen-Specker theorem [1] asserts that, in a Hilbert space with a finite number of dimensions, $d \geq 3$, it is possible to produce a set of n projection operators, representing yes-no questions about a quantum system, such that none of the 2^n possible answers is compatible with the sum rules of quantum mechanics. Namely, if a subset of mutually orthogonal projection operators sums up to the unit matrix, one and only one of the answers is yes. The physical meaning of this theorem is that there is no way of introducing noncontextual “hidden” variables [2] which would ascribe definite outcomes to these n yes-no tests. This conclusion holds irrespective of the quantum state of the system being tested.

It is also possible to formulate a “state-specific” version of this theorem, valid for systems which have been prepared in a known pure state. In that case, the projection operators are chosen in a way adapted to the known state. A smaller number of questions is then sufficient to obtain incompatibility with the quantum mechanical sum rules. An even smaller number is needed if strict sum rules are replaced by weaker probabilistic arguments [3,4].

The original proof by Kochen and Specker [1] involved projection operators over 117 vectors in a 3-dimensional real Hilbert space $\mathbb{R}^3$. A simple proof with 33 vectors was later given by Peres [5], who also reported an unpublished construction by Conway and Kochen, using only 31 vectors [6]. A proof with 20 vectors in $\mathbb{R}^4$ was recently given by Kernaghan [7]. Here, we consider the Kochen-Specker theorem in $\mathbb{R}^8$. A state-independent proof is produced with 36 vectors, which can be collected into 30 distinct projection operators. A state-specific proof involves only 13 vectors, and achieves the lowest value of the ratio n/d that has been obtained so far.

Our construction is based on Mermin’s remark [8] that, for any three spin- $\frac{1}{2}$ particles, the four operators

$$A = \sigma_{1z} \otimes \sigma_{2z} \otimes \sigma_{3z}, \tag{1}$$

$$B = \sigma_{1z} \otimes \sigma_{2x} \otimes \sigma_{3x}, \tag{2}$$

$$C = \sigma_{1x} \otimes \sigma_{2z} \otimes \sigma_{3x}, \tag{3}$$

$$D = \sigma_{1x} \otimes \sigma_{2x} \otimes \sigma_{3z}, \tag{4}$$

commute. Moreover, their product is proportional to the unit matrix,

$$ABCD = -\mathbf{1}. \tag{5}$$

Mermin's three particle system is the simplest one that illustrates *both* Bell's theorems, on nonlocality [9] and contextuality [10].

Each one of the five equations above involves a complete set of commuting operators, and therefore determines a complete orthogonal basis in $\mathbb{R}^8$ (complex numbers are not needed here, because σ_x and σ_z are real matrices). The five orthogonal octads generated by the above equations are listed in the first column of Table 1. To simplify typography, each vector was given norm 2 (this avoids the use of fractions) and the symbol $\bar{1}$ means -1 . The components of each vector are written as a horizontal array, rather than the usual "column vector." The basis used is the direct product of the bases where each σ_z is diagonal. For example, the vector with components 00002000 is the eigenvector for which σ_{1z} has eigenvalue -1 , and σ_{2z} and σ_{3z} have eigenvalue 1. This is most easily seen by using binary digits, 0 and 1, for labelling the "up" and "down" components of a spinor, respectively, and combining them into binary numbers, 000, 001, ..., 111, for labelling components of vectors in $\mathbb{R}^8$. Thus, the vector 00002000 has the physical meaning stated above because its only nonvanishing component is the 100th one (that is, the fifth one, in binary notation).

It is easy to see that the 40 projection operators on these 40 vectors provide an example of the Kochen-Specker contradiction. Indeed, associating values 0 (no) and 1 (yes) to the vectors in a basis (with a single 1, of course) amounts to selecting one of these eigenvectors, and the latter indicates which ones of the four commuting operators which generate that basis have value 1, and which ones have value -1 . Such a mapping of each operator on one of its eigenvalues cannot be done for all of them in Eqs. (1–5): it would lead to an inconsistency, because of the minus sign in (5). This argument is readily generalized to a larger number of spin- $\frac{1}{2}$ particles, and provides a proof of the Kochen-Specker theorem in a real Hilbert space with 2^n dimensions. However, fewer than 2^n vectors are actually needed for the proof, as we shall see.

First, we note that the above set of 40 vectors has a high degree of symmetry. (Unfortunately, we have not been able to determine its invariance group. We asked several well known experts, who also did not find it.) A direct inspection, best done by computer, shows that each vector is orthogonal to 23 other ones (7 in the same basis, and 4 in each one of the four other bases), and it makes a 60° or 120° angle with each one of the 16 remaining vectors. It is possible to construct with the 40 vectors 25 distinct orthogonal octads (each vector appears in 5 octads). Eleven of these octads are listed in the remaining columns of Table 1 (the 14 other octads are not needed for the proof and have not been listed).

It is seen that there are four vectors, namely those with components 20000000, 00001111, 001 $\bar{1}$ 001 $\bar{1}$, and 10 $\bar{1}$ 0 $\bar{1}$ 010, that do not appear in the 11 octads (they appear of course in the 14 unlisted octads). These four vectors are mutually orthogonal, and they belong to four different of the original octads. It is easily seen that there are 1280 different ways of choosing four vectors with these properties (that is, of choosing which are the 11 octads, out of 25, that appear in Table 1).

It is also seen that each one of the 36 remaining vectors appears in Table 1 either twice or four times. Let us now consider the projection operators over these 36 vectors. According to quantum mechanics, each projection operator corresponds to a yes-no question: the eigenvalues 1 and 0 mean yes and no, respectively. Each orthogonal octad defines 8 commuting projection operators (that is, 8 compatible questions) which sum up to the unit matrix. This means that if an experimental test is actually performed for these 8 questions, the answer is yes to one, and only one of them. The value 1 is thereby associated with one of the eight vectors of each octad, and the value 0 with the others.

If nothing is known of the state of the system, quantum mechanics is unable to predict which vector will get the value 1. A natural question is whether there could be a more complete theory, such that the value associated with each vector would be determined by “hidden variables.” Table 1 readily shows that this goal cannot be achieved. Indeed, the sum of values in each octad is always 1, therefore the sum of values for the 11 octads in the table is 11, which is an *odd* number. On the other hand, each vector (tentatively associated with a value which is either 0 or 1) appears either twice or four times in the table (with the *same* value), thus contributing 2 or 4 (an *even* number) to the sum of values. We have reached a contradiction. This is the proof of the Kochen-Specker theorem in $\mathbb{R}^8$.

Furthermore, our 36 incompatible propositions can be combined into a smaller number, namely 30 distinct ones, by using projection operators of rank 2 on the *planes* spanned by some pairs of vectors. Table 2 shows how 20 of the vectors can be combined into 14 planes. The vectors on each line are mutually orthogonal. Each plane is spanned by two adjacent vectors on the same line (for example, 02000000 and 00000002). When such a pair of vectors occurs in any of the 11 octads listed in Table 1 (that is, in one of the columns) this vector pair should be replaced by the corresponding plane. Once this is done, each column lists planes and unpaired vectors, all of which are mutually orthogonal. They correspond to commuting projection operators of rank 1 or 2, which sum up to **11**. Therefore, if “experimental” (that is, counterfactual) values 0 or 1 are attributed to them, these values sum up to 11, exactly as before. On the other hand, each one of the remaining 16 vectors (those not used in Table 2) appears twice in the new version of Table 1, and

each of the 14 planes appears 2 or 4 times – always an even number. We thus reach the same Kochen-Specker contradiction as before.

Until now, we assumed nothing about the state of the quantum system. If that state is known with certainty, it becomes possible to find a much smaller number of propositions which lead to a Kochen-Specker contradiction. For instance, let the quantum system be prepared in the pure state $100\bar{1}0\bar{1}\bar{1}0$ (this does not restrict the generality of this discussion in any way, because it is always possible to choose the basis in $\mathbb{R}^8$ in such a way that any given pure state be represented by a vector with these components). We can now discard from Table 1 that vector, whose associated value is 1, by definition, and all the vectors orthogonal to it, whose associated values are zero. Only 7 non-empty columns, with 13 different vectors, remain in Table 1. In each column, there are four mutually orthogonal vectors, which span a subspace containing the known pure state $100\bar{1}0\bar{1}\bar{1}0$ (because the complementary orthogonal subspace is also orthogonal to $100\bar{1}0\bar{1}\bar{1}0$). Explicitly, we have

$$2 \times 100\bar{1}0\bar{1}\bar{1}0 = 1010\bar{1}0\bar{1}0 + 10\bar{1}010\bar{1}0 - 00020000 - 00000200, \quad (6)$$

$$= 1100\bar{1}\bar{1}00 + 1\bar{1}001\bar{1}00 - 00020000 - 00000020, \quad (7)$$

$$= 11\bar{1}\bar{1}0000 + 1\bar{1}1\bar{1}0000 - 00000200 - 00000020, \quad (8)$$

$$= 11\bar{1}\bar{1}0000 + 00001\bar{1}\bar{1}1 + 1010\bar{1}0\bar{1}0 - 01010101, \quad (9)$$

$$= 1\bar{1}1\bar{1}0000 + 00001\bar{1}\bar{1}1 + 1100\bar{1}\bar{1}00 - 00110011, \quad (10)$$

$$= 1100\bar{1}\bar{1}00 + 1\bar{1}001\bar{1}00 + 001\bar{1}00\bar{1}1 - 00110011, \quad (11)$$

$$= 1100\bar{1}\bar{1}00 + 001\bar{1}00\bar{1}1 + 10\bar{1}010\bar{1}0 - 01010101. \quad (12)$$

Quantum mechanics asserts that, for each one of the above equations, there exists an experimental test which randomly selects one of the four vectors appearing on the right hand side of that equation. A hidden variable theory would claim that the selection is not random, and that there are preassigned values, 0 and 1, corresponding to all 13 vectors: on each line, one of the four vectors has value 1, the others have value 0. Moreover, we may demand that, if the same vector appears in different lines, the value associated to it is everywhere the same (that is, the result of an experimental test which would determine that value is not “contextual”). These demands then lead us to the same contradiction: there are 7 equations, so that the sum of values is 7. On the other hand, each vector appears 2 or 4 times, so that the sum of values is even. It is remarkable that no more

than 13 propositions are needed to reach the contradiction. The proof entirely holds in these seven equations.

Work by AP was supported by the Gerard Swope Fund, and the Fund for Encouragement of Research.

1. S. Kochen and E. P. Specker, J. Math. Mech. 17 (1967) 59.
2. M. Redhead, Incompleteness, nonlocality, and realism (Clarendon Press, Oxford, 1987).
3. R. Clifton, Am. J. Phys. 61 (1993) 443.
4. H. Bechmann Johansen, Am. J. Phys. 62 (1994) 471.
5. A. Peres, J. Phys. A: Math. Gen. 24 (1994) L175.
6. A. Peres, Quantum theory: concepts and methods (Kluwer, Dordrecht, 1993) p. 114.
7. M. Kernaghan, J. Phys. A: Math. Gen. 27 (1994) L829.
8. N. D. Mermin, Phys. Rev. Lett. 65 (1990) 3373; Rev. Mod. Phys. 65 (1993) 803.
9. J. S. Bell, Physics 1 (1964) 195.
10. J. S. Bell, Rev. Mod. Phys. 38 (1966) 447.

Table 1. Orthogonal octads used for proving the theorem.

20000000									
02000000	×							×	
00200000	×						×		
00020000							×	×	
00002000	×					×			
00000200						×		×	
00000020						×	×		
00000002	×					×	×	×	
11110000		×				×			
11 $\bar{1}\bar{1}$ 0000						×			×
1 $\bar{1}\bar{1}\bar{1}$ 0000						×		×	
1 $\bar{1}\bar{1}$ 10000	×					×		×	×
00001111									
000011 $\bar{1}\bar{1}$	×								×
00001 $\bar{1}\bar{1}\bar{1}$	×							×	
00001 $\bar{1}\bar{1}$ 1								×	×
11001100							×	×	
1100 $\bar{1}\bar{1}$ 00		×					×	×	×
1 $\bar{1}$ 001 $\bar{1}$ 00		×					×		
1 $\bar{1}$ 00 $\bar{1}$ 100							×		×
00110011		×						×	
001100 $\bar{1}\bar{1}$								×	×
001 $\bar{1}$ 001 $\bar{1}$									
001 $\bar{1}$ 00 $\bar{1}$ 1		×							×
10101010			×					×	×
1010 $\bar{1}$ 0 $\bar{1}$ 0								×	×
10 $\bar{1}$ 010 $\bar{1}$ 0								×	×
10 $\bar{1}$ 0 $\bar{1}$ 010									
01010101			×					×	
01010 $\bar{1}$ 0 $\bar{1}$									×
010 $\bar{1}$ 010 $\bar{1}$			×						×
010 $\bar{1}$ 0 $\bar{1}$ 01			×						×
100101 $\bar{1}$ 0	×	×	×	×					
100 $\bar{1}$ 0110	×	×	×		×				
10010 $\bar{1}$ 10	×				×				
100 $\bar{1}$ 0 $\bar{1}\bar{1}$ 0	×	×		×	×				
0110 $\bar{1}$ 001				×	×				
01 $\bar{1}$ 01001		×	×	×	×				
0 $\bar{1}$ 101001		×			×				
0 $\bar{1}\bar{1}$ 0 $\bar{1}$ 001			×		×				

Table 2. Construction of 14 planes from 20 vectors.

02000000	00000002	00000020	
11110000	1 $\bar{1}\bar{1}$ 10000	00001 $\bar{1}\bar{1}$ 1	
1 $\bar{1}$ 001 $\bar{1}$ 00	1100 $\bar{1}\bar{1}$ 00	001100 $\bar{1}\bar{1}$	
01010101	10101010	01010 $\bar{1}$ 0 $\bar{1}$	
10010 $\bar{1}$ 10	100101 $\bar{1}$ 0	01 $\bar{1}$ 01001	0 $\bar{1}$ 101001
0 $\bar{1}$ $\bar{1}$ 0 $\bar{1}$ 001	100 $\bar{1}$ 0110	100 $\bar{1}$ 0 $\bar{1}\bar{1}$ 0	0110 $\bar{1}$ 001